

Action of Diosgenin and homoeopathic pathogenesis of *Dioscorea villosa*

Kisor Kumar Naskar^{1*}, Ompriya Mishra²

¹Department of Materia Medica, D. N. De Homoeopathic Medical College and Hospital, ²Department of Obstetrics and Gynaecology, National Institute of Homoeopathy, Kolkata, West Bengal, India

Abstract

Background: *Dioscorea villosa* is a popular herbal remedy for pains. Early Americans used this to treat colic which is also established in homoeopathic proving and verified clinically along with other important phenomena. Diosgenin is its primary active ingredient. It has been investigated that all possible revealed actions of this plant are due to this ingredient. **Objective:** The aim of this study is to understand pathological background and prescribing areas of *D. villosa* by correlating and analyzing Homoeopathic pathogenesis with the knowledge of pathophysiological activities of Diosgenin. **Study Method:** Information was collected from various researches on animal and human models regarding actions of Diosgenin. Homoeopathic pathogenesis of *D. villosa* have been studied and analyzed from 'The Guiding Symptoms of our Materia Medica' by C. Hering and also from the experiences of different authors to find pathological background as well as prescribing areas of *D. villosa* by correlating with the actions of Diosgenin. **Result:** Diosgenin has prominent action on cholesterol metabolism, hormones (oestrogen and dehydroepiandrosterone [DHEA]), inflammations, and allergic activities on animal and human models. These pathological actions of Diosgenin have similarity with the pathological background of homoeopathic pathogenesis of *D. villosa*. **Conclusion:** Actions of Diosgenin reflect in the homoeopathic pathogenesis of *D. villosa* such as altered cholesterol metabolism, irregularity of hormonal (DHEA and estrogen) functions, and anti-inflammatory and anti-allergic phenomena with their related manifestations. Moreover, these might be the underlying pathological essence of this drug.

Keywords: Anti-allergic, Anti-cancerous, Anti-diabetic, Anti-inflammatory, Cholesterol metabolism, Dehydroepiandrosterone-oestrogen, *Dioscorea villosa*, Diosgenin, Homoeopathic pathogenesis

INTRODUCTION

Dioscorea villosa (Wild Yam) is named after the Greek Physician Dioscorides of the 1st century. It contains saponins, namely Dioscin, which gives Diosgenin on hydrolysis. It is the precursor for synthesis of steroidal hormones. It also contains Dioscorin as an alkaloid.

Diosgenin is considered as the primary active ingredient. Dioscin has similar or overlapping effects to Diosgenin, but there has been little investigation done on the effects of Dioscin alone, Dioscorin has no known physiologic effects. Diosgenin is structurally similar to cholesterol.^[1]

Due to structural similarity, its action on cholesterol metabolism is not only hypothesized but has also been proven in several research studies. Diosgenin promotes fecal cholesterol excretion by inhibiting intestinal cholesterol absorption and enhancing biliary secretion.^[2]

Diosgenin has oestrogen properties. It may bind with oestrogen receptors of hypothalamus and takes part in negative feedback mechanism of estrogen control.^[3]

Wild yam is considered as a precursor to dehydroepiandrosterone (DHEA). DHEA supplements are manufactured from it and considered as a valid source, but the exact mechanisms of its conversion toward hormones *in vivo* are unknown. Some research studies proved its activity suggesting the same.^[4-6]

In several studies conducted in animal and human models, it has significant anti-inflammatory, anti-allergic, anti-diabetic, anti-cancerous properties as well.

***Address for correspondence:** Dr. Kisor Kumar Naskar, 393, Dhamaitala School Road, Dakshin Jagaddal, Rajpur Sonarpur, Kolkata - 700 151, West Bengal, India. E-mail: drkisor_naskar@yahoo.co.in

Access this article online

Quick Response Code:



Website:
www.ijrh.org

DOI:
10.4103/0974-7168.200845

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Naskar KK, Mishra O. Action of diosgenin and homoeopathic pathogenesis of *Dioscorea villosa*. Indian J Res Homoeopathy 2017;11:5-11.

ACTION OF DIOSGENIN

Action in cholesterol metabolism

From several studies, it has been found that Diosgenin has the following properties related to cholesterol metabolism.

In animal models, it has been found that it inhibits cholesterol absorption.^[7,8] Side by side, it increases biliary cholesterol secretion and output as well as altered serum cholesterol levels,^[8-10] increases fecal excretion of cholesterol, decreases plasma cholesterol levels,^[11,12] decreases low-density lipoprotein (LDL) cholesterol, and elevates high-density lipoprotein (HDL) cholesterol.^[13] Not only that, it may also protect from bile salt toxicity.^[14] In human model, it has been found that Diosgenin can decrease triglycerides and phospholipids but increase serum HDL.^[15] In another study in mice model, Diosgenin significantly decreased serum total cholesterol level while slightly increased HDL level and also found protective effect against atherosclerosis by regulating pro-inflammatory mediators in serum.^[16]

Action on oestrogen and dehydroepiandrosterone

Although there is no such direct evidence of conversion of Diosgenin into any hormone, several studies suggest its oestrogen property.^[17] This property can be established by its binding with estrogen receptors of the hypothalamus as a part of negative feedback mechanism of estrogen control.

It significantly increased DHEA level in another study on rats.^[18]

Action on inflammations and allergic activities

It has significant anti-inflammatory activity as found in different studies in animal models as on sub-inflammatory intestinal inflammation in indomethacin-induced intestinal inflammation in rats,^[19] showing immunosuppressive effects by reducing the production of inflammatory mediators.^[20] It also has regulatory action on T-cell immunity in intestine of mice with food allergy,^[21] suppressive action of immunoglobulin E production on allergen-induced intestine inflammation, and mast cell infiltration and degranulation of food allergy in a murine model.^[22] It is also found that Diosgenin has significant anti-inflammatory activity by the inhibition of lipopolysaccharide-induced pro-inflammatory cytokines in a dose-dependent manner without any cytotoxicity^[23] and proved useful to ameliorate the inflammatory changes in obese adipose tissues.^[24]

Action in diabetes

It has anti-diabetic property involving glucose metabolism and might be a novel therapeutic agent.^[25,26] In a study, it has been found that Diosgenin-induced increased DHEA may contribute to the improvement of hyperglycemia through muscular glucose transporter Type 4 signaling pathway activation in type 1 diabetes of rats.^[18]

Action in cancer

Different studies also revealed the action of Diosgenin on cancer. Several evidences suggest that it can reduce tumor

multiplicity of adenoma or adenocarcinoma in colon,^[27,28] breast adenocarcinoma,^[29,30] hepatocellular carcinoma,^[31,32] gastric cancer,^[33] prostate cancer,^[34] osteosarcoma,^[35] etc.

HOMOEOPATHIC PATHOGENESIS OF DIOSCOREA VILLOSA

Important pathological conditions as well as manifestations of *D. villosa* from detailed recorded pathogenesis are seen in 'The Guiding Symptoms of our Materia Medica' by C. Hering^[36] and from the clinical description based on the experiences of pioneers like Dr. T.F. Allen,^[37] Dr. William H. Burt,^[38] Dr. J.C. Burnett,^[39] Dr. Samuel Lilienthal,^[40] Dr. P.N. Verma and Dr. Indu Vaid^[41] and others.

According to the proving of Dr. Gushing, there was dual action of *Dioscorea*. In primary effects as erotism and in secondary effects as impotency and both are attended by its peculiar type of spermatorrhea. Dr. Boyd reported that *Dioscorea* has curative property in spasmodic stricture of urethra with "pain about the umbilicus relieved by pressure; pressure on the rectum; and paroxysmal colic pains."^[42]

Gastrointestinal system

- Colic - Flatulent colic, bilious colic, and neuralgia of bowels with spasmodic pain in abdomen. Colic with feeble digestive power relieved by stretching the body out or by walking about. This modality has been considered as a characteristic feature of *Dioscorea*
- Diarrhoea and dysentery with colic
- Liver affection – Hard, dry, black stool indicates a diminution of bile, but large doses produce increased secretion of bile as indicated by the papaceous, yellow, loose stool^[38]
- Gall stone and much flatulence;^[40] along with feeble digestion.^[41]

Reproductive system

- Increased sexual excitement - constant sexual excitement, nocturnal emissions with amorous dreams, and erections when asleep, sometimes without erections. There is also diminished sexual desire with no erections for many days.
- Severe pain with cramp in flexor tendons of fingers and toes, alternating with false labor pains.

Vascular affections

- Haemorrhoids with constipation.^[38] Haemorrhoids with darting pain shooting to liver. It looks like bunches of grapes with great pain and distress^[37]
- Acute painful varicocele from excess venery or long-lasting unsatisfied desire^[39]

Heart affections

Angina pectoris, pains radiate into both arms.

Extremities

- Felons - early, when pains are sharp
- Sciatica right sided, shooting down thigh, <motion.

ASSOCIATING HOMOEOPATHIC PATHOGENESIS TO ACTIONS OF DIOSGENIN

Effects of altered cholesterol metabolism

- Colic - Symptoms of biliary colic with flatulence and its character has underlying relation with the property of altered cholesterol metabolism
- Modality of colic - Pain while reclining and relieved by sitting up which is often retroperitoneal^[43]
- Pancreatic cause of abdominal colic - Pancreatic cause of abdominal colic and its affections can also be related to secondary cause of altered cholesterol metabolism or estrogenic effects^[44-46]
- Atherosclerotic affection of heart - Heart symptoms because of atherosclerotic changes in the arteries, which is due to the same altered cholesterol metabolism. It is found that Diosgenin has protective effect against atherosclerosis^[16]
- Atherosclerosis and sciatica - Atherosclerosis of arteries supplying the lumbar region has been suggested as a mechanism leading to inter-vertebral disc degeneration causing sciatica^[47]
- Hyperlipidemia and neuralgias. In a study, it has been identified that hyperlipidemia is a novel risk factor for peripheral neuropathies. It includes axonal distal polyneuropathy, vision and hearing loss, motor nerve system lesions, and sympathetic nervous system dysfunction.^[48]

Irregularities of hormonal functions

- Sexual excitement, vivid dreams, and emissions - In sexual sphere, action of DHEA can be related to vivid dreams and emissions. However, in a recent study, it has been found that reducing estrogen in men undergoing treatment for low testosterone resulted in a drop in libido and an increase in body fat, even as testosterone levels improved^[49]
- False labor-like pain - Oestrogen has a property to induce the formation of oxytocin receptors in the uterus and make uterus sensitive to oxytocin for expulsion of fetus. Moreover, it also established that oxytocin receptor expression is being modulated by stimulation with estrogen not only in the pregnant but also in the nonpregnant uterus. Hence, estrogenic activity of *Dioscorea* not only facilitates labor by increasing oxytocin receptors density in uterus but it is also responsible for uterine contractility-associated disorders of the nonpregnant uterus^[50]
- Constipation - Oestrogen has a significant role to develop constipation^[51]
- Chronic venous diseases - In a recent study, it was also found that estrogen receptors play an important role in chronic venous diseases^[52]
- Urethral stricture - In case of male, a large amount of estrogen cause atrophy and fibrosis of the spongy and cavernous bodies of the penis, depriving the urethra of its normal elasticity, which results in stricture.^[53]

Anti-allergic and anti-inflammatory actions

- Diarrhoea or dysentery with colic - Abdominal colic with bowel complaints related to Diarrhoea or dysentery might be related to food allergy^[22]
- Panaritium or felon - Symptoms of peripheral inflammatory changes in the form of panaritium might be related to the anti-inflammatory property of Diosgenin. It can also be explained from the pathology of hypercholesterolemia. It has been found that hypercholesterolemia is also associated with lymphatic vascular changes, leads to obstruction of lymphatic drainage. As a result of that, lack of protective body immune response may maintain this infection as well as inflammatory process.^[54]

PRESCRIBING AREAS REVEALED FROM ANALYZED HOMOEOPATHIC PATHOGENESIS AND ALSO FROM DIFFERENT ACTIONS OF DIOSGENIN

1. Biliary, flatulent, or neurogenic colic of abdomen
2. Gall stone
3. Intestinal allergic or inflammatory affections with diarrhea, dysentery, and colic
4. Hypercholesterolemia or dyslipidemia and related manifestations, including atherosclerotic affections of heart and vascular system, peripheral vascular affection such as claudication
5. Complaints or manifestations related to DHEA and estrogen hormonal irregularities including sexual erotism or impotency with spermatorrhea
6. Oestrogen effects on venous system such as constipation with hemorrhoids, varicocele
7. Urethral stricture
8. Neuralgia
9. Diabetes
10. Cancer - Colon tumors (adenoma-adenocarcinoma); breast adenocarcinoma (HER2 oncoprotein-overexpressing AU565 human [also in MCF-7 and MDA-231]); hepatocellular cancer; gastric cancer; prostate cancer; and in osteosarcoma.^[27-35]

Co-existence of manifestations derived from or associated with pathological essence of *Dioscorea* such as altered cholesterol metabolism, irregular hormonal (estrogen-DHEA) actions, allergic and inflammatory activities are the basis of homoeopathic approach as logical combination to prescribe *Dioscorea*. Similarly, characteristic pains or sensations, its extension, modalities, or other characteristic phenomena such as sexual dreams and spermatorrhea as stated by pioneers may represent individualizing features along with these underlying pathologies.

CONCLUSION

As expression and intensity of disease manifestations differ from one to another, drug pathogenesis revealed from homoeopathic

proving and clinical experiments also differ because of individual's constitutional tendencies and susceptibility. However, inspite of these differing manifestations, the target areas and tendencies of the drug are definite. Similarly, drug actions on general or local physiological activities with altered function of internal factor(s) including hormone, enzyme, salt, or other bio-chemicals are also definite. In other words, varying degrees of intensity and differing modalities, sensations, extensions, or other modifications of resulting manifestations are developed from the definite altered physiological activities with affections of different bio-chemicals. These pathophysiological changes with altered function of bio-chemical or other factors are the pathological essence of medicine.

From the present study of *Dioscorea*, it can be identified that effect of pathological activity of Diosgenin shown in animal as well as in human models has similarity in the internal pathophysiological changes in homoeopathic pathogenesis of *D. villosa*. External signs and symptoms of these internal changes were expressed fragmentarily with individualistic differences throughout the recorded pathogenesis of this homoeopathic medicine.

Hence, it can be concluded that the underlying altered cholesterol metabolism, irregularity of hormonal (DHEA and estrogen) function, and anti-inflammatory and anti-allergic phenomena with their related manifestations are reflected as the pathological essence of *Dioscorea*. And in the similar way of producing these pathological changes, *Dioscorea* has a capability to influence individual's homoeostatic mechanism to maintain the normal state of these altered pathology and related phenomena.

In addition to the pathological complaints related to hypercholesterolemia or dyslipidemia, irregularity of estrogen and DHEA, future evaluation in the sphere of malignancies in colon, breast, hepatic, gastric, prostate, bones as well as for diabetes and intestinal allergic affections are to be considered for more clinical applicability of *Dioscorea* in homoeopathic practice.

Financial support and sponsorship

Nil.

Conflict of interest

None declared.

REFERENCES

1. Anjali K, Kathi JK. Wild Yam (Dioscoreaceae); The Longwood Herbal Task Force. Available from: <http://www.mcp.edu/herbal/default.htm>. [Last revised 1999 Jun 23].
2. Temel RE, Brown JM, Ma Y, Tang W, Rudel LL, Ioannou YA, et al. Diosgenin stimulation of fecal cholesterol excretion in mice is not NPC1L1 dependent. *J Lipid Res* 2009;50:915-23.
3. Michelle M. Herbs for the treatment of the symptoms of menopause – A phytotherapist's perspective. Available from: www.mediherb.com/media/pp-no-51. [Last revised 2011 May].
4. Rector-Page LG. Healthy Healing – A Guide to Self-healing for Everyone. 11th ed. United States of America: Traditional Wisdom Inc.; 2000. p. 90.
5. Fuchs NK. The Health Detectives' 456 Most Powerful Healing

Secrets: Easy Read edi. Accessible Publishing System PTY Ltd – ReadHowYouWant. Aus. 2008. p. 241.

6. Elkins R. DHEA. Pleasant Grove, Utah: Woodland Publishing; 1996. p. 6.
7. Juarez-Oropeza MA, Diaz-Zagoya JC, Rabinowitz JL. *In vivo* and *in vitro* studies of hypocholesterolemic effects of diosgenin in rats. *Int J Biochem* 1987;19:679-83.
8. Nervi F, Bronfman M, Allalón W, Depiereux E, Del Pozo R. Regulation of biliary cholesterol secretion in the rat. Role of hepatic cholesterol esterification. *J Clin Invest* 1984;74:2226-37.
9. Cayen MN, Dvornik D. Effect of diosgenin on lipid metabolism in rats. *J Lipid Res* 1979;20:162-74.
10. Nervi F, Marinovic I, Rigotti A, Ulloa N. Regulation of biliary cholesterol secretion. Functional relationship between the canalicular and sinusoidal cholesterol secretory pathways in the rat. *J Clin Invest* 1988;82:1818-25.
11. Uchida K, Takase H, Nomura Y, Takeda K, Takeuchi N, Ishikawa Y. Changes in biliary and fecal bile acids in mice after treatments with diosgenin and beta-sitosterol. *J Lipid Res* 1984;25:236-45.
12. Malinow MR, Elliott WH, McLaughlin P, Upson B. Effects of synthetic glycosides on steroid balance in *Macaca fascicularis*. *J Lipid Res* 1987;28:1-9.
13. Cayen MN, Dvornik D. Combined effects of clofibrate and diosgenin on cholesterol metabolism in rats. *Atherosclerosis* 1978;29:317-27.
14. Accatino L, Pizarro M, Solis N, Koenig CS. Effects of diosgenin, a plant-derived steroid, on bile secretion and hepatocellular cholestasis induced by estrogens in the rat. *Hepatology* 1998;28:129-40.
15. Araghiniknam M, Chung S, Nelson-White T, Eskelson C, Watson RR. Antioxidant activity of dioscorea and dehydroepiandrosterone (DHEA) in older humans. *Life Sci* 1996;59:PL147-57.
16. Park HJ, Byeon HE, Koo HJ, Pyo S. Anti-atherosclerotic effects of diosgenin in ApoE-deficient mice. *FASEB J* 2011; 25:980.2
17. Aradhana, Rao AR, Kale RK. Diosgenin – A growth stimulator of mammary gland of ovariectomized mouse. *Indian J Exp Biol* 1992;30:367-70.
18. Sato K, Fujita S, Iemitsu M. Diosgenin administration improve hyperglycemia through the activation of muscular glucose metabolism. *FASEB J* 2013;27:1154.26.
19. Yamada T, Hoshino M, Hayakawa T, Ohhara H, Yamada H, Nakazawa T, et al. Dietary diosgenin attenuates subacute intestinal inflammation associated with indomethacin in rats. *Am J Physiol* 1997;273:G355-64.
20. Jung DH, Park HJ, Byun HE, Park YM, Kim TW, Kim BO, et al. Diosgenin inhibits macrophage-derived inflammatory mediators through downregulation of CK2, JNK, NF-kappaB and AP-1 activation. *Int Immunopharmacol* 2010;10:1047-54.
21. Huang CH, Liu DZ, Jan TR. Diosgenin, a plant-derived sapogenin, enhances regulatory T-cell immunity in the intestine of mice with food allergy. *J Nat Prod* 2010;73:1033-7.
22. Huang CH, Ku CY, Jan TR. Diosgenin attenuates allergen-induced intestinal inflammation and IgE production in a murine model of food allergy. *Planta Med* 2009;75:1300-5.
23. Singh M, Hamid AA, Maurya AK, Prakash O, Khan F, Kumar A, et al. Synthesis of diosgenin analogues as potential anti-inflammatory agents. *J Steroid Biochem Mol Biol* 2014;143:323-33.
24. Hirai S, Uemura T, Mizoguchi N, Lee JY, Taketani K, Nakano Y, et al. Diosgenin attenuates inflammatory changes in the interaction between adipocytes and macrophages. *Mol Nutr Food Res* 2010;54:797-804.
25. McAnuff MA, Omoruyi FO, Morrison EY, Asemota HN. Changes in some liver enzymes in streptozotocin-induced diabetic rats fed sapogenin extract from bitter yam (*Dioscorea polygonoides*) or commercial diosgenin. *West Indian Med J* 2005;54:97-101.
26. Ghosh S, More P, Derle A, Patil AB, Markad P, Asok A, et al. Diosgenin from *Dioscorea bulbifera*: Novel hit for treatment of type II diabetes mellitus with inhibitory activity against α -amylase and α -glucosidase. *PLoS One* 2014;9:e106039.
27. Miyoshi N, Nagasawa T, Mabuchi R, Yasui Y, Wakabayashi K, Tanaka T, et al. Chemoprevention of azoxymethane/dextran sodium sulfate-induced mouse colon carcinogenesis by freeze-dried yam sanyaku and its constituent diosgenin. *Cancer Prev Res (Phila)* 2011;4:924-34.

28. Raju J, Bird RP. Diosgenin, a naturally occurring steroid [corrected] saponin suppresses 3-hydroxy-3-methylglutaryl CoA reductase expression and induces apoptosis in HCT-116 human colon carcinoma cells. *Cancer Lett* 2007;255:194-204.
29. Chiang CT, Way TD, Tsai SJ, Lin JK. Diosgenin, a naturally occurring steroid, suppresses fatty acid synthase expression in HER2-overexpressing breast cancer cells through modulating Akt, mTOR and JNK phosphorylation. *FEBS Lett* 2007;581:5735-42.
30. Srinivasan S, Koduru S, Kumar R, Venguswamy G, Kyprianou N, Damodaran C. Diosgenin targets Akt-mediated prosurvival signaling in human breast cancer cells. *Int J Cancer* 2009;125:961-7.
31. Li F, Fernandez PP, Rajendran P, Hui KM, Sethi G. Diosgenin, a steroidal saponin, inhibits STAT3 signaling pathway leading to suppression of proliferation and chemosensitization of human hepatocellular carcinoma cells. *Cancer Lett* 2010;292:197-207.
32. Li Y, Wang X, Cheng S, Du J, Deng Z, Zhang Y, *et al.* Diosgenin induces G2/M cell cycle arrest and apoptosis in human hepatocellular carcinoma cells. *Oncol Rep* 2015;33:693-8.
33. Mao ZJ, Tang QJ, Zhang CA, Qin ZF, Pang B, Wei PK, *et al.* Anti-proliferation and anti-invasion effects of diosgenin on gastric cancer BGC-823 cells with HIF-1a shRNAs. *Int J Mol Sci* 2012;13:6521-33.
34. Chen PS, Shih YW, Huang HC, Cheng HW. Diosgenin, a steroidal saponin, inhibits migration and invasion of human prostate cancer PC-3 cells by reducing matrix metalloproteinases expression. *PLoS One* 2011;6:e20164.
35. Moalic S, Liagre B, Corbière C, Bianchi A, Dauça M, Bordji K, *et al.* A plant steroid, diosgenin, induces apoptosis, cell cycle arrest and COX activity in osteosarcoma cells. *FEBS Lett* 2001;506:225-30.
36. Hering C. *The Guiding Symptoms of our Materia Medica*. Vol. 5. New Delhi: B. Jain Pub. Pvt. Ltd.; 2000. p. 125-42.
37. Allen TF. *A Hand Book of Materia Medica and Homoeopathic Therapeutics*. Kolkata: New Central Book Agency Pvt. Ltd.; 2007. p. 459.
38. William HB. *Physiological Materia Medica*. New Delhi: B. Jain Pub. (P) Ltd.; 2000. p. 382-3.
39. Chitkara HL. *Best of Burnett*. New Delhi: B. Jain Pub. (P) Ltd.; 2010. p. 182.
40. Samuel L. *Homoeopathic Therapeutics*. New Delhi: B. Jain Pub. Pvt. Ltd.; 1996. p. 199.
41. Verma PN, Indu V. *Encyclopaedia of Homoeopathic Pharmacopoeia and Drug Index*. Vol. II. 4th ed., New Delhi: B. Jain Pub. (P) Ltd.; 2007. p. 986.
42. Hale EM. *Dioscorea – Its Action on Genital Organs; Hahnemannian Monthly*. 1868-69. p. 306. Available from: <http://www.homeopathyworldcommunity.com>. [Posted on 2010 Oct 4]
43. Charles H. *Abdominal Pain; Department of Family Medicine – OSU College of Osteopathic Medicine*. Available from: docplayer.net/11013584-Abdominal-pain-charles-henley-do-mph-department-of-fam. [Last revised on 2002 Sep].
44. Dominguez-Muñoz JE, Malfertheiner P, Ditschuneit HH, Blanco-Chavez J, Uhl W, Büchler M, *et al.* Hyperlipidemia in acute pancreatitis. Relationship with etiology, onset, and severity of the disease. *Int J Pancreatol* 1991;10:261-7.
45. Yadav D, Pitchumoni CS. Issues in hyperlipidemic pancreatitis. *J Clin Gastroenterol* 2003;36:54-62.
46. Rodríguez Santana Y, Nimo Román A, García Sáez I, López Alvarez JM, Consuegra Llapur E, González Jorge R. Treatment of severe hypertriglyceridemia with continuous insulin infusion. *Case Rep Crit Care* 2011;2011:293917.
47. Leino-Arjas P, Kauppila L, Kaila-Kangas L, Shiri R, Heistaro S, Heliövaara M. Serum lipids in relation to sciatica among Finns. *Atherosclerosis* 2008;197:43-9.
48. Song Wu, Xu Cao, Rongzhen He, Kun Xiong. Detrimental impact of hyperlipidemia on the peripheral nervous system - A novel target of medical epidemiological and fundamental research study; *Neural Regen Res* 2012.
49. Finkelstein JS, Lee H, Burnett-Bowie SA, Pallas JC, Yu EW, Borges LF, *et al.* Gonadal steroids and body composition, strength, and sexual function in men. *N Engl J Med* 2013;369:1011-22.
50. Richter ON, Kübler K, Schmolling J, Kupka M, Reinsberg J, Ulrich U, *et al.* Oxytocin receptor gene expression of estrogen-stimulated human myometrium in extracorporeally perfused non-pregnant uteri. *Mol Hum Reprod* 2004;10(5):339-46
51. Oh JE, Kim YW, Park SY, Kim JY. Estrogen rather than progesterone cause constipation in both female and male mice. *Korean J Physiol Pharmacol* 2013;17:423-6.
52. Serra R, Gallelli L, Perri P, De Francesco EM, Rigracciolo DC, Mastroberroto P, *et al.* Estrogen receptors and chronic venous disease. *Eur J Vasc Endovasc Surg* 2016;52:114-8.
53. Yokoyama M, Fukutani K, Kawamura T, Shoji F, Ohtani M. Stricture of the anterior urethra following estrogen therapy in patients with prostatic cancer. *Urol Int* 1983;38:247-50.
54. Lim HY, Rutkowski JM, Helft J, Reddy ST, Swartz MA, Randolph GJ, *et al.* Hypercholesterolemic mice exhibit lymphatic vessel dysfunction and degeneration. *Am J Pathol* 2009;175:1328-37.

1. Die Wirkung von Diosgenin und die homöopathische Pathogenese von *Dioscorea villosa*

Auszug

Hintergrund: *Dioscorea villosa* ist ein beliebtes pflanzliches Schmerzmittel. Früher verwendeten die Amerikaner wilden Yams zur Behandlung von Koliken. Es wurde auch eine Arzneiprüfung durchgeführt und durch wichtige Phänomene klinisch verifiziert. Diosgenin ist der Hauptwirkstoff. Es wurde untersucht, ob alle möglichen Wirkungen dieser Pflanze auf diesen Wirkstoff zurückzuführen sind.

Ziel: Es ist eine Studie zum Verständnis des pathologischen Hintergrunds und der Verordnung von *Dioscorea villosa* durch Korrelation und Analyse homöopathischer Pathogenesen mit dem Wissen über pathophysiologische Aktivitäten von Diosgenin.

Methoden: Es wurden Informationen aus verschiedenen Forschungen über tierische und menschliche Modelle zur Wirkung von Diosgenin gesammelt. Die homöopathische Pathogenese von *Dioscorea villosa* wurde anhand von C. Herings "The Guiding Symptoms of Our Materia Medica" und Erfahrungen verschiedener Autoren untersucht und analysiert, um pathologische Ursachen zu finden und Anwendungsbereiche von *Dioscorea villosa* in Korrelation zur Wirkung von Diosgenin zu beschreiben.

Ergebnisse: Diosgenin hat eine auffällige Wirkung auf den Cholesterinstoffwechsel, auf Hormone [Estrogen und DHEA (Dehydroepiandrosteron)], Entzündungen und allergische Aktivitäten durch seine pathologischen Veränderungen bei tierischen und menschlichen Modellen. Diese pathologischen Wirkungen von Diosgenin haben Ähnlichkeit mit dem pathologischen Hintergrund homöopathischer Pathogenesen von *Dioscorea villosa*.

Fazit: Die Wirkungen von Diosgenin spiegeln sich in den homöopathischen Pathogenesen der *Dioscorea villosa* als veränderter Cholesterinstoffwechsel, in Unregelmäßigkeiten der hormonellen (DHEA- und Östrogen-) Funktionen, bei entzündungshemmenden und antiallergischen Phänomene mit ihren verwandten Manifestationen wider, Und dies könnte die pathologische Hauptursache dieser Droge sein.

Acción de la Diosgenina y Patogenesis Homeopática de *Dioscorea Villosa*

RESUMEN

Fundamentos: La *Dioscorea villosa* es un medicamento vegetal frecuentemente utilizado en el dolor. Los primeros americanos recurrían al ñame silvestre para tratar cólicos, lo cual también se ha establecido en las patogenesis homeopáticas y se ha verificado clínicamente, junto con otros fenómenos importantes. La diosgenina es su principal ingrediente. Se ha podido verificar que todas las posibles acciones de esta planta se deben a este componente.

Objetivos: Este estudio se efectuó para conocer la base patológica y las indicaciones de *Dioscorea villosa* correlacionando y analizando las patogenesis homeopáticas con el conocimiento de las actividades fisiopatológicas de la diosgenina.

Método de estudio: La información se recopiló a partir de diferentes investigaciones en modelos animales y humanos sobre las acciones de la diosgenina. Se estudiaron y analizaron las patogenesis homeopáticas de *Dioscorea villosa* de "The Guiding Symptoms of our Materia Medica" de C. Hering, así como de otras experiencias de diferentes autores para determinar las bases patológicas y las indicaciones de *Dioscorea villosa* en correlación con las acciones de la diosgenina.

Resultados: La diosgenina posee efectos prominentes en el metabolismo del colesterol, las hormonas (estrógeno y DHEA [dehidroepiandrosterona]), así como en las actividades inflamatorias y alérgicas a través de los cambios patológicos en los modelos animales y humanos. Estas acciones patológicas de la diosgenina son similares a la patología observada en las patogenesis homeopáticas de *Dioscorea villosa*.

Conclusiones: Las acciones de la diosgenina quedan reflejadas en las patogenesis homeopáticas de *Dioscorea villosa* como un metabolismo alterado del colesterol, en irregularidades en las funciones hormonales (DHEA y estrógenos), y en los fenómenos antiinflamatorios y antialérgicos con sus manifestaciones relacionadas. Esto puede ser la esencia patológica subyacente de este fármaco.

डाईओसजेनिन के प्रभाव और डायस्कोरिया विल्लोसा की रोगजनन क्षमता

सार

पृष्ठभूमि: डायस्कोरिया विल्लोसा दर्द के लिए एक लोकप्रिय हर्बल उपचार है। प्रारंभिक अमेरिकियों द्वारा जंगली रतालू का उपयोग पेट दर्द (उदरशूल) के उपचार हेतु किया गया, जिसे होम्योपैथिक औषध प्रमाणन तथा नैदानिक सत्यापन द्वारा भी स्थापित किया गया है। डायोसजेनिन इसका प्राथमिक सक्रिय संघटक है। यह जाँचा गया है कि इस पादप के संभाव्य प्रभाव इस घटक के कारण ही हैं।

उद्देश्य: डायोसजेनिन की पैथो-फिजिओलॉजिकल गतिविधियों के ज्ञान के साथ होम्योपैथिक रोगजनन के विश्लेषण और सहसंबंध द्वारा डायस्कोरिया विल्लोसा की रोगजनन पृष्ठभूमि और निर्धारित क्षेत्रों की समझ के लिए एक अध्ययन।

अध्ययन विधि: जानवरों और मानव मॉडल पर डायोसजेनिन के प्रभाव से संबंधित विभिन्न अनुसंधानों से प्राप्त जानकारीयों एकत्रित की गईं। डायस्कोरिया विल्लोसा की होम्योपैथिक रोगजनन क्षमता का अध्ययन किया गया तथा सी. हरिंग द्वारा लिखित 'द गाइडिंग सिम्पटम्स ऑफ अवर मेटोरिया मेडिका' से विश्लेषण किया गया। डायस्कोरिया विल्लोसा की रोगजनन पृष्ठभूमि व विस्तार क्षेत्र का अध्ययन डायोसजेनिन के प्रभावों के सहसंबंध द्वारा किया गया।

परिणाम: डायोसजेनिन का जानवर और मानव मॉडलों पर अपने रोग परिवर्तन के दौरान कोलेस्ट्रॉल उपापचय, हार्मोन्स (इस्ट्रोजन और डीएचइए (डिहाइड्रोएपिएन्ड्रोस्टेरोन)), सूजन और एलर्जी की गतिविधियों पर प्रमुख प्रभाव है। डायोसजेनिन के इन रोगजनन प्रभावों की डायस्कोरिया विल्लोसा की होम्योपैथिक रोगजनन की पृष्ठभूमि के साथ समानता पाई गई।

निष्कर्ष: डायोसजेनिन के प्रभाव डायस्कोरिया विल्लोसा की होम्योपैथिक रोगजनन क्षमता में दर्शित होते हैं जैसे बदला हुआ कोलेस्ट्रॉल उपापचय, हार्मोनल अनियमितता (डीएचइए और ईस्ट्रोजन), सूजन विरोधी और एलर्जी विरोधी प्रतिभास। यह इस औषधि का अंतर्निहित रोग सार हो सकता है।