

Erratum: A randomized comparative trial in the management of Alcohol Dependence: individualized Homoeopathy versus standard Allopathic treatment

In the article titled, “A randomized comparative trial in the management of Alcohol Dependence: individualized Homoeopathy versus standard Allopathic treatment” published in pages 172-181, issue 3, vol. 10 of Indian Journal of Research in Homoeopathy,^[1] the first sentence under the section “Results” in the abstract of the journal is written incorrectly as “ITT analysis reflected 80% ($n = 32$) of the patients in IH and 37.5% ($n = 15$) of the patients in the SA responding to CI before 2.4 treatment with absolute difference was 42.5% (42.5 [95% confidence interval [CI]: 23.0, 61.6]) and estimated effect: 6.6 (95% C.I: 2.4, 18.2), $P = 0.0002$.”

The correct sentence should be read as:

“ITT analysis reflected 80% ($n = 32$) of the patients in IH and 37.5% ($n = 15$) of the patients in the SA responding to treatment with absolute difference of 42.5% (95% confidence interval [CI]: 23.0, 61.6) and estimated effect: 6.6 (95% CI: 2.4, 18.2, $P = 0.0002$).”.

REFERENCE

1. Manchanda RK, Janardanan Nair KR, Varanasi R, Oberai P, Bhuvanewari R, Bhalerao R, *et al*. A randomized comparative trial in the management of Alcohol Dependence: Individualized Homoeopathy versus standard Allopathic Treatment. Indian J Res Homoeopathy 2016;10:172-81.

DOI: 10.4103/0974-7168.200840