

## CLINICAL RESEARCH

### A Prospective Multicentric Observational Study to Evolve the Usefulness of the Predefined Homoeopathic Medicines in the Management of Acute Diarrheal Disease in Children\*\*

C. Nayak<sup>1\*</sup>, Vikram Singh<sup>1</sup>, K. Singh<sup>1</sup>, Hari Singh<sup>1</sup>, Anita Sharma<sup>1</sup>, Praveen Oberai<sup>1</sup>, Varanasi Roja<sup>1</sup>, Debadatta Nayak<sup>1</sup>, Maya Padmanabhan<sup>1</sup>, Alok Mishra<sup>1</sup>, G.C. Sehegal<sup>2</sup>, Subhash Kaushik<sup>2</sup>, C.P. Chowdhary<sup>2</sup>, V.K. Singh<sup>3</sup>, Sahid Ali<sup>4</sup>, P. Hima Bindu<sup>4</sup>, D.B. Sarkar<sup>5</sup>

1. Central Council for Research in Homoeopathy, Headquarters, New Delhi, India
2. Central Research Institute, Noida (Uttar Pradesh), India
3. Homeopathic Drug Research Institute, Lucknow (Uttar Pradesh), India
4. Drug Standardization Unit, extension centre, Hyderabad (Andhra Pradesh)
5. Drug Proving Research Unit, Kolkata (West Bengal)

**Background & objectives:** Acute diarrheal disease is one of the major causes of morbidity and mortality in developing countries. No data regarding the pediatric population suffering from acute diarrhea seeking homoeopathic treatment in India are available. Therefore, an observational study on acute diarrheal diseases in children was undertaken with 14 predefined trial homoeopathic medicines to arrive at a group of useful homeopathic medicines for the treatment of acute childhood diarrhea.

**Methods:** A prospective, multi-centre observational study was conducted by Central Council for Research in Homeopathy at its various Institutes and Units throughout India during Oct. 2005- Sept. 2008. A total of three hundred twenty seven children in the age-group, 6 months to 12 years, were included. Trial medicines selected on the principles of Homeopathy were prescribed and the diarrhea index score was assessed before and after treatment using SPSS (ver. 16).

**Results:** The difference in the mean number of stools and diarrhea index score was found to be statistically significant ( $p=0.000$ ,  $<0.05$ ) after the prescription of trial homoeopathic medicines in 321 children. The diarrhea index score got either worsened or remained unchanged from the baseline score in 6 (1.83%) children. Of the 14 medicines half of them viz., *Podophyllum peltatum* ( $n=158$ ), *Chamomilla* ( $n=49$ ), *Aethusa cynapium* ( $n=25$ ), *Mercurius solubilis* ( $n=23$ ), *Calcarea carbonica* ( $n=21$ ), *Sulphur* ( $n=18$ ), and *Phosphorus* ( $n=17$ ) were found to be most useful among the 14 predefined trial medicines.

**Conclusion:** This was an observational study with positive results leads to further validation by suitable Randomized Control Trial(s).

**Key words:** homeopathy; acute diarrheal disease in children; podophyllum peltatum; chamomilla

\*\*This article was originally published in the journal, American Journal of Homoeopathic Medicine, Vol. 102, No. 3, Autumn 2009. Reprint with the consent of the Authors and the Publisher.

\* Address for correspondence

Prof. C.Nayak

Director General

Central Council for Research in Homoeopathy

61-65, Institutional area, opp. D Block

Janakpuri, New Delhi, India

Pin code- 110058

Email-ccrh@del3.vsnl.net.in

## Introduction

Acute diarrheal disease is one of the major causes of morbidity and mortality in the developing world, especially among infants and children below 5 years of age [1]. Estimates suggest that during the 1990s, nearly 1.4 billion diarrhea episodes occurred every year among children younger than 5 years of age in socio-economically developing countries, of which 123.6 million episodes required outpatient medical care and 9 million episodes required hospitalization. Approximately 2 to 2.5 million diarrhea-associated deaths were estimated annually in this age group, concentrated in the most impoverished areas of the world [2]. The percentage of children under 3 years of age who suffered from diarrhea in the two-week period before the survey was 10% in the National Family Health Survey (NFHS) – 1 and 19% in NFHS-2 [3, 4]. Estimations of the burden of diarrhoeal diseases in India by the National Institute of Cholera and Enteric Diseases (NICED), Kolkata indicates that diarrhoeal diseases contribute to about 9.1% of deaths in the age group of 0–6 years. It has been further estimated that in India the years of life lost (YLL) due to diarrhoeal diseases in the 0-6 years' population presently contribute to about 98% of disability adjusted life-years (DALYs) and would probably remain unchanged over the next decade till 2016 [5]. For children under 5 years of age in developing areas and countries, there was a median of 3.2 episodes of diarrhea per child-year [6]. This indicated little change from previously described incidences. There has not been a concurrent decrease in morbidity rates attributable to diarrhea. As population growth is focused in the poorest areas, the total morbidity component of the disease burden is greater than previously [6].

The homeopathic system of therapy, an outcome of experimental research, is based on the principle of "Similia Similibus curenter" which implies that a drug cures in the sick what it causes in the healthy. According to World Health Organization, Homeopathy is the second most used health care system in the world [7]. It has shown its effectiveness in treating both acute as well as chronic illnesses [8]. Homeopathy helped in controlling cholera epidemic [9] where the mortality rate was 8.5% in comparison to 51.5% in allopathic treatment group. Three randomized clinical trials [RCT] and one meta-analysis of these trials of homeopathic medicines on acute childhood diarrhea by Jacobs *et al* [10-13] proved to be statistically significant in comparison to the placebo group.

In homeopathy, the treatment being person-specific and not disease-specific, there is invariably more than one useful medicine for any disease condition. As such, feasibility of RCT in homeopathy

is still an enigma. Moreover, before conducting an RCT we need to know about the population intended for study. No data regarding the pediatric population suffering from acute diarrhea seeking for homeopathic treatment in India are available. So the present observational study on acute diarrheal diseases in children was undertaken with 14 predefined trial medicines to arrive at a group of efficacious homeopathic medicines for the treatment of acute childhood diarrhea.

## Objectives

### Primary objective

To evolve a group of efficacious homeopathic medicines in the management of acute diarrheal disease in children.

### Secondary objectives

- i. To verify characteristic symptoms of medicines used.
- ii. To ascertain clinical symptoms, if any, in respect of homeopathic medicines used during the study.

## Methods

### Study design

The study was a prospective, multi-centre observational study conducted at various Institutes/Units: Central Research Institute (CRI), Noida (Uttar Pradesh), Homeopathic Drug Research Institute (HDRI), Lucknow (Uttar Pradesh), Drug Proving Research Unit (DPRU), Kolkata (West Bengal) and Drug Standardization Unit (DSU), extension centre, Hyderabad (Andhra Pradesh) under Central Council for Research in Homeopathy (henceforth Council) during the period October 2005- September 2008. A diarrhea index score (S1) was designed to assess the severity of the illness (Table 1). A second score (S2) from 1-7 was designed by the Council, thus a child reporting for treatment on the first day of infection was given highest score of 7 and one reporting with seven days of infection, the lowest score, 1. The study protocol was in accordance with the Helsinki declaration [14] on human experimentation. Necessary ethical clearance was obtained from Council's Ethical Committee.

### Patient population

Four hundred and sixty nine children, between 6 months and 12 years of age, with 3 unformed stools for at least 2 consecutive days, were eligible to

**Table 1:** Diarrhea index score

Symptom/signs	Score				
	0	1	2	3	4
No. of Stools / day	Normal	4	6	8	12
Diarrhoea (consistency)	Normal stool	Loose	Watery	Slimy	Mucus
Fever	Absent		Low grade		High grade
Vomiting / day	Absent	Only once	2 – 3	3 – 5	
Gross/Occult blood in stool	Absent			Fresh blood	Black
Weakness	Absent	Mild	More		Profound has to rest
Muscular cramps	Absent	Mild	Moderate	Severe	
Weight loss	None			< 5%	Mild (5-6%)

participate. Each child was followed up to recovery or maximum of 7 days. Children reporting with diarrhea of prolonged duration of more than 7 days, symptoms suggesting cholera, associated high grade fever, pseudo diarrhea or frequent passage of small volumes of stool of normal consistency, fecal incontinence (for reasons other than diarrhea), parasitic/protozoa infestations, salmonella organisms causing gastroenteritis, persistent diarrhea, children on any other medicine for diarrhea, bloody stool were excluded. Diarrhea with moderate and severe dehydration as per WHO guidelines were also excluded [15].

#### Selection of medicine

The selection of trial medicines was done by repertorising the nosological symptoms [16] of acute diarrhea. Tyler M & Weir J's elimination method [17] of repertorization was used for selection of trial medicines. 'Diarrhoea Children in' was taken as the eliminating rubric and thus 14 medicines of 3 marks (first grade) were short listed using Complete Repertory in Cara professional software [18]. These medicines were: *Aethusa cynapium*, *Calcarea sulph.*, *Calcarea carbonica*, *Chamomilla*, *Magnesia muriatica*, *Mercurius solubilis*, *Psorinum*, *Ipecacuanha*, *Rheum palmatum*, *Silicea*, *Stramonium*, *Sulphur*, *Phosphorus* and *Podophyllum peltatum*. These medicines were procured from M/s Sharda Boiron laboratory Pvt. Ltd., Sahibabad, India. Each child received a homeopathic medicine selected on the basis of presenting totality. However, if the choice of medicine was outside of the trial medicines then that child was not enrolled and treated in general out patient department.

#### Potency, doses & repetition

All the enrolled patients were given medicine in 6c (10<sup>-12</sup> dilution) potency in frequent doses (2-6 hourly) depending upon frequency, duration and intensity of

the symptoms. Each dose consisted of 4 globules, size no. 30. After the first prescription, if improvement remains stand still next higher potency i.e. 30c was given. Placebo globules were given as soon as the improvement was observed.

#### Treatment plan

Homeopathic treatment was given as per instructions given in Hahnemann's Organon of Medicine [19]. Its characteristics are: Selection of one drug at a time, using the 'similia principle' and the drug picture and disease picture should be as similar as possible. Investigators were allowed to change the prescription upto three times from among the trial medicines if the first prescription didn't work. As a part of non-medical management all the guardians/parents were advised to make their child take oral rehydration therapy (ORS), if necessary.

#### Outcome assessment

A diarrhea index score was used to quantify the severity of illness as mentioned in the study design. The diarrhea index score (S1) was divided according to intensity of the disease into mild (2-7), moderate (8-18) and severe (19-26). Assessment was done by calculating the percentage using the formula [(S1 at baseline - S1 at completion of the study) / S1 at baseline]. Thus 100% was defined as cure, >75% - <100% as marked, 50% - <75% as moderate, 25% - <50% as mild, < 25% as insignificant improvement, 0% as not improved or status quo and any increase in symptom score from the baseline score was counted as worse.

#### Statistical analysis

Descriptive statistical characteristics and comparative analysis like t test and one way Anova was done using SPSS (Statistical package for social

science) Version 16. A linear regression model was used to adjust for disparities in patient characteristics at the initial visit.

## Results

During the two years eleven months period, 469 children were enrolled by different Institutes/Units under the Council, out of which 25 were lost to follow up, 14 were withdrawn, 103 were excluded due to their enrolment on 1<sup>st</sup> & 2<sup>nd</sup> day of illness, however these

children were followed up and the data were analyzed separately. A total of 327 children (194 males; 133 females) as per the inclusion and exclusion criteria were followed up and studied [Mean age of children less than 1 year was 10.1±2.9 months; children above 1 year was 5.6±3.2 years]. The incidence of acute diarrhea was mostly in 1-5 years of age group (n=155) followed by 6-10 years age group (n=125), and < 1 year (n=47). Descriptive characteristics of children at baseline are given in Table 2 & 3.

**Table 2:** Baseline details of children suffering from acute diarrhea

	Study group (n)	Mean ±SD	%
Institutes/Units			
CRI, Noida	114		34.9
HDRI, Lucknow	107		32.7
DSU (extension), Hyderabad	68		20.8
DPRU, Kolkata	38		11.6
Sex			
• Male	194		59.3
• Female	133		40.7
Day of infection			
• 3 <sup>rd</sup>	265		81
• 4 <sup>th</sup>	54	2.67±0.95	16.5
• 5 <sup>th</sup>	8		2.4
Mean days of diarrhea	327	3.19±0.4	100
Diarrhoea index score (range)			
• Mild (2-7)	218	5.2±1.3	66.6
• Moderate (8-18)	109	9.2±1.3	33.4
No. of stools per day	327	7.2±2.7	100

**Table 3:** Presenting symptoms / signs of acute diarrhea in children

Symptoms/signs (n)	At baseline			At end		
	No. of children	Percentage (%)	Total %	No. of children	Percentage (%)	Total %
No. of stools per day 4/6/8/12	69/125/68/65	20.8/38.2/21.1/19.9	100	6/2/2/1	1.8/0.6/0.6/0.3	3.3
Diarrhoea (consistency) Loose/watery/ slimy/mucoid	51/248/26/2	15.6/75.8/8/0.6	100	9/5/2/1	2.7/1.5/0.6/0.3	5
Fever Low grade/high grade	144/1	43.9/0.3	44.2	7/1	2.1/0.3	0
Vomiting/day Only once/ 2-3 times/3-5 times	73/59/16	22.3/18/4.9	45.2	3/2/1	0.9/0.6/0.3	1.8
Weakness Mild/more/ profound has to rest	92/26/5	28.1/8/1.5	37.6	¼/2	0.3/1.2/0.6	2.1
Muscular cramps Mild	4	1.2	1.2	4	1.2	1.2
Weight loss (dehydration) Less than 5%	9	2.7	49.7	9	2.7	2.7

The S1 was evaluated at baseline and at end of the study using paired t test (Table 4). The difference in the mean S1 was found to be statistically significant (P= 0.000, <0.05). A paired t test for the average no. of stools per day for all children indicated a statistically significant difference between baseline and at the end of the treatment (t=44.2, df =326, P= 0.000, <0.05).

A linear regression model containing the variables age, sex, diarrhea- index score, number of stools in the past 48 hours was used (because age, diarrhea index score, number of stools were found to be highly

correlated, only the variable age was used in the model as independent factor with no. of stools at end as the dependent factor). The result of this analysis is statistically significant (p =0.000, <0.05).

An analysis of the trial medicines (Table 5) showed that 7 trial medicines which were used frequently in prescribing to children (n=312, 95.1%) suffering from acute diarrhoea are *Podophyllum peltatum* (n=158; 48.3%), *Chamomilla* (n=49, 15%), *Aethusa cynapium* (n=25;7.6%), *Calcarea carbonica* (n=21,6.4%), *Mercurius solubilis* (n=23, 7%), *Sulphur*(n=18, 5.5%),

**Table 4:** Diarrhoea index score at entry and at end

	Mean score at entry ± SD (n)	Mean score at end ± SD (n)	95% confidence interval level
Diarrhea index score	6.5±2.2 (327)	0.3±1.7 (327)*	5.94-6.52
Range in diarrhea index score			
• Mild (2-7)	5.3±1.3 (218)	0.2±1.3 (218)*	4.75-5.26
• Moderate (8-18)	9.2±1.3 (109)	0.5±2.1 (109)*	8.27-9.11
No. of stools per day	7.2±2.7 (327)	0.2±1.1 (327)*	6.68-7.30

\*Statistically significant using paired t test, p-value = 0.000, <0.05

**Table 5:** Data of trial medicines in the study

Medicine	No. of patients	Percent	p-value	95% confidence interval difference	Outcome assessment <sup>†</sup>					
					Cured	Marked >	Mode rate>	Mild >	<	=
Podophyllum	158	48.3	0.000	5.65-6.46	149	7	1	0	1	0
Chamomilla	49	15	0.000	6.08-7.38	48	0	1	0	0	0
Aethusa cynapium	25	7.6	0.000	5.98-8.73	22	0	1	1	1	0
Mercurius solubilis	23	7	0.000	5.12-7.48	20	1	1	0	1	0
Calcarea carbonica	21	6.4	0.000	3.95-6.62	20	0	0	0	1	0
Sulphur	18	5.5	0.000	5.11-7.22	18	0	0	0	0	0
Phosphorus	17	5.2	0.000	5.21-7.84	16	1	0	0	0	0
Ipecacuanha	7	2.1	0.000	4.55-9.44	7	0	0	0	0	0
Magnesium muriaticum	2	0.6			0	0	0	0	0	2
Psorinum	2	0.6	0.042	1.14-13.85	1	1	0	0	0	0
Rheum	2	0.6	0.182*	-23.36-40.26	2	0	0	0	0	0
Silicea	2	0.6	0.126*	-3.85-8.85	1	0	0	1	0	0
Calcarea sulphurica	1	0.3	0.000		1	0	0	0	0	0
Stramonium	0	0	0	0	0	0	0	0	0	0
Total	327				305	10	4	2	4	2

\*Not significant, p=> 0.05; <sup>†</sup>>- Improvement; < worse; = status quo

and *Phosphorus* (n=17, 5.2%). Among the trial medicines 4.9% (n=6) less used in this study are *Ipecacuanha* (n=7), *Magnesia muriatica* (n=2), *Psorinum* (n=2), *Rheum palmatum* (n=2), *Silicea* (n=2), *Calcarea sulphurica* (n=1). The characteristic indications of the medicines found useful in 10 or more children are described in Table 6.

Data of children who were enrolled within 48 hours of suffering from diarrhea are as follows: 103 (59 males; 44 females); [Mean age of children less than 1 year was 9.6±2.2 months; children above 1 year were 4.7±3 years]. The mean no. of stools of these children at entry and at end was 7.2±2.7 and 0.1±1.1 respectively. Paired t test was done related to diarrhea index score and the results were found to be statistically significant (p=0.000, <0.05).

One way ANOVA was carried out for all the children (n=431) making three groups according to days of infection (1<sup>st</sup> – 2<sup>nd</sup>, 3<sup>rd</sup>, and 4<sup>th</sup> -5<sup>th</sup> day of illness) and their improvement status respectively. It was found that children belonging to 1<sup>st</sup>-2<sup>nd</sup> day of infection responded more in comparison to other two groups, F (2, 429)=8.64, p=0.000 <0.05.

93.2% of children (n=305) were cured at various days of their treatment (Figure 1). Most of the children (n=109) were cured on 2<sup>nd</sup> day of their treatment. Only

10 children had showed marked degree of improvement and 4 children mild improvement while 4 children got worsening of their diarrhea and were referred for other treatment and 2 children remained status quo.




Figure 1: Trend of cure in children at different days of illness and treatment

### Discussion

The present study is one of its kinds in India for treating acute diarrhea in children with homeopathic medicines and reflects the positive results of homeopathic therapy in children suffering from acute diarrhea. This study also supports the findings of the previous studies[10-12] that individualized

Table 6: Characteristic indications of frequently used medicines

Name of trial medicine	Indications
<i>Podophyllum peltatum</i>	Diarrhea painless, putrid, with prostration, < hot weather, < morning; stool profuse, forcible with noise; profuse thirst. Color of stool changes with each stool.
<i>Chamomilla</i>	Irritable, wants to be carried, cries during sleep. Stool is watery, greenish; very offensive, smells like rotten eggs, <evening, night. Thirsty, drinks eagerly. Dentitional diarrhea.
<i>Aethusa cynapium</i>	Complete absence of thirst; weakness & prostration. Milk intolerance, can't digest milk/ intolerance to milk. Sleepiness after each stool. Diarrhea during dentition.
<i>Calcarea carbonica</i>	Slow in activities; fatty and flabby children, profuse sweating esp. from head; obstinate. Sour smelling stool, white chalky stool. Milk intolerance.
<i>Mercurius solubilis</i>	Hyperactive, destructive, disobedient. Profuse thirst with moist tongue. Slimy offensive stool, excoriation around anus, tenesmus before and after stool, < night. Profuse sweating, too offensive.
<i>Phosphorus</i>	Chilly patient; lean, thin, affectionate child. Longs for cold drinks. Stool contains sago-like particles; great weakness and prostration. Vomiting <after drinking,
<i>Sulphur</i>	Craving for sweet; aversion to bath. Anal orifice red; empty sensation in stomach; early morning diarrhea, stool contains undigested food; burning sensation in anus. Diarrhea associated with skin eruptions.

homeopathic treatment decreases the duration of diarrhea and number of stools in children with acute diarrhea.

However, there was no data regarding the pediatric population suffering from acute diarrhea coming for homeopathic treatment in India. Moreover, India is a country otherwise known as sub-continent with diverse geographical, cultural and dietary variations. This made us to take this study in observational design, although RCTs [10-12] were conducted in the past outside India, but in limited population.

Children with diarrhea of 3-7 days duration were enrolled; some of them may be on the way of natural recovery. But most of the patients (80.8%) enrolled on 3<sup>rd</sup> day of their illness and only (2.4%) were enrolled on 5<sup>th</sup> day of illness. Moreover, according to Strina *et al* [20] acute diarrhea may persist till 8-13<sup>th</sup> day, as 4% of children fall in this category. So bias due to natural regression of disease might be negligible.

In this study we observed that the mean no. of stool at entry and at end was (7.2±2.7; 0.2±1.1). The result showed statistically significant decrease in the no. of stool (p=0.000) and thus was similar to the findings of Jacob's [10-12] (7.9±3.6 at baseline)[11]. Similarly mean days of suffering from diarrhea in our study was found to be 3.2±0.4 and the days required for diarrhea index score to become zero is 3.2±1.5 which further corroborates with findings of Jacob *et al* [11]. But the difference is that in this study we enrolled children after two consecutive days of unformed stool while the later enrolled the children with 3 unformed stools within 24 hours. On considering all the cases (n=430) it was observed that mean days of suffering from diarrhea at entry was 2.9±0.6 and the mean days for diarrhea index score to become zero was 3.5±1.5, which is less than Jacob *et al* [12] findings (3.6±1.9).

Similar to previous studies [11, 12] male children were more affected than female children. This study was conducted both in urban and suburban areas. Though the children were advised to report daily till 7<sup>th</sup> day of illness but this couldn't be achieved because as soon as the child starts improving the parent reported irregularly. So data for all the days of treatment was not available though the final recovery was reported.

The trial medicines useful in this observational study were *Podophyllum*, *Chamomilla*, *Mercurius solubilis*, *Aethusa cynapium*, *Phosphorus* and *Calcarea carbonica* which are similar to the medicines used by Jacobs *et al* [12], but *Arsenicum album* was found to be useful in the latter study whereas this medicine was not used in our study as it was not considered, because it is mentioned in second degree [18].

Stool of all the children at the screening level were examined for protozoa, parasite and if found these patients were excluded. Stool culture was not done to ascertain microbiological cause of diarrhea. But in one patient the cyst for *Entamoeba histolytica* was found when the stool was re-examined on worsening, the same lacked at enrollment level. Such data limits the analysis of the effect of homeopathic medicines on pathogens causing diarrhea which is a limitation of this study.

The strength of this study is that it represents a pragmatic setting of homeopathic practice which reflects the day-to-day clinical practice. To ascertain clinical symptoms in respect of homeopathic medicines was one of the objectives of the study. But such symptoms were not observed which re-confirms the completeness of these medicines which are in use since centuries.

As the study did not have any control group, randomization and blinding, it can't conclude that homeopathic therapy is effective in acute diarrhea prevalent among the children of India. To validate the effect of homeopathic care on acute diarrhea further research is warranted in same population. Effect of homeopathic medicine on acute diarrhea of specific etiology should be considered in future studies.

## Conclusion

This was an observational study with positive results and these results need further validation by suitable Randomized Control Trial(s).

## Acknowledgements

We thank Dr. Abha Aggarawal, Deputy Director, National Institute of Medical Statistics, New Delhi for reviewing the statistical data. Programme officers of respective Institutes/Units where the study was conducted are acknowledged for administrative support.

## Conflict of interest

We declare no conflict of interest.

## References

1. Snyder DC, Merson MH. The magnitude of the global problem of acute diarrhoeal disease: A review of active surveillance data. *Bull. World Health Organ.* 60 (4) 605-613 (1982)
2. Ryan Miguel O, Prado Valeria, and Pickering Larry K., A Millennium Update on Pediatric Diarrheal Illness in the Developing World; *Semin Pediatr Infect Dis* 2005; 16:125-136

3. International Institute for Population Sciences (IIPS) and ORC Macro. 2000. *National Family Health Survey (NFHS-2), 1998–99: India*. Mumbai: IIPS
4. International Institute for Population Sciences (IIPS). 1995. *National Family Health Survey (MCH and Family planning), India 1992–1993*: Bombay: IIPS.
5. Burden of Disease in India National Commission on Macroeconomics and Health Ministry of Health & Family Welfare, Government of India, New Delhi September 2005
6. Kosek Margaret, Bern Caryn, & Guerrant Richard L. The global burden of diarrhoeal disease, as estimated from studies published between 1992 and 2000; *Bull. World Health Organ.* 2003; 81:197-204.
7. Chapman Edward; Homeopathy. In Jonas W.B., Levin J.S., Essentials of Complementary Medicine; 1999; Lipincott Williams Wilkins [CD ROM].
8. Kleijnen J, Knipschild P, Ter Riet G. Clinical trials of Homeopathy, *BMJ* ; 1991; 302; 316-23
9. Gaucher Catherine, Jeulin Dominique, Peycru Patrick, Pla Anna, Amengual Carles. Cholera and Homeopathic medicine The Peruvian experience, *British Homeopathic Journal* 1993 (82):155-163.
10. Jacobs Jennifer, Jimenez Luz Margarita, Gloyd Steven, Carares Felix Esponzia, Gaitan Nargarita Paniagua, Crothers Dean. Homeopathic treatment of acute childhood diarrhea A randomized clinical trial in Nicaragua. *British Homeopathic Journal* 1993 (82):83-86)
11. Jacobs Jennifer, Jimenez L. Margarita, Gloyd S Stephen, Gale L.J., Crothers D. Treatment of Acute Childhood Diarrhea with homeopathic medicine: A randomized clinical trial in Nicaragua. *Pediatrics*; vol. 93 No. 5 May 1994, pp. 719-725.
12. Jacobs Jennifer, L. Margarita Jimenez, Malthouse Stephen, Chapman Elizabeth, Crothers Dean, Masuk Mary, and Jonas Wayne B., Homeopathic Treatment of Acute Childhood Diarrhea: Results from a Clinical Trial in Nepal; *The Journal Of Alternative And Complementary Medicine*; 6 (2), 2000 : 131-139)
13. Jacobs Jennifer, Jonas Wayne B., Jimenez L. Margarita, Crothers D. Homeopathy for childhood diarrhea: combined results and metaanalysis from three randomized clinical trials. *Pediatr Infect Dis J*; 2003; 22: 229-34.
14. World Medical Association Declaration of Helsinki -Ethical Principles for Medical Research Involving Human Subjects. Available from URL: [http://www.who.int/bulletin/archives/79\(4\)373.pdf](http://www.who.int/bulletin/archives/79(4)373.pdf) , accessed on 8.8.2005.
15. World Health Organization. *Readings on diarrhoea: student manual*. Geneva, World Health Organization, 1992
16. Limbos Mary Ann. Approach to the child with diarrhea. In Osborn Lucy M, Dewitt Thomas G, First Lewis R, Zenel Joseph A. editors *Pediatrics*, 2005: p-628
17. Tyler M.L., Weir J. Repertorising In Kent J.T. *Repertory of the Homeopathic Materia Medica*; Reprint 6<sup>th</sup> American Edition; New Delhi; B Jain Publishers; 1989
18. Witko, David. *CARA Professional* © 1997, London, Miccant Ltd. Revised programme by John Stevenson. 1999
19. Haahnemann Samuel. *Organon of Medicine*; 5<sup>th</sup> & 6<sup>th</sup> edition; New Delhi; B Jain Publishers; 1994
20. Strina Agostino, Cairncross Sandy, Prado Matildes S., Teles Carlos A.S., Barreto Mauricio L. Childhood diarrhea symptoms, management and duration: observations from a longitudinal community study. *Transactions of the Royal society of Tropical Medicine and Hygiene* (2005) 99, 407-416